

**Patient Information**  
**MAVYRET™ (MAV-ih-reht)**  
(glecaprevir and pibrentasvir)  
Tablets

**What is the most important information I should know about MAVYRET?**

**MAVYRET can cause serious side effects, including:**

**Hepatitis B virus reactivation.** Before starting treatment with MAVYRET, your healthcare provider will do blood tests to check for hepatitis B virus infection. If you have ever had hepatitis B virus infection, the hepatitis B virus could become active again during or after treatment of hepatitis C virus with MAVYRET. Hepatitis B virus becoming active again (called reactivation) may cause serious liver problems including liver failure and death. Your healthcare provider will monitor you if you are at risk for hepatitis B virus reactivation during treatment and after you stop taking MAVYRET.

**For more information about side effects, see the section “What are the possible side effects of MAVYRET?”**

**What is MAVYRET?**

- MAVYRET is a prescription medicine used to treat adults with chronic (lasting a long time) hepatitis C virus (HCV) genotypes 1, 2, 3, 4, 5 or 6 infection without cirrhosis or with compensated cirrhosis.

MAVYRET contains the two medicines: glecaprevir and pibrentasvir.

It is not known if MAVYRET is safe and effective in children under 18 years of age.

**Do not take MAVYRET if you:**

- have certain liver problems
- also take any of the following medicines:
  - atazanavir (EVOTAZ<sup>®</sup>, REYATAZ<sup>®</sup>)
  - rifampin (RIFADIN<sup>®</sup>, RIFAMATE<sup>®</sup>, RIFATER<sup>®</sup>, RIMACTANE<sup>®</sup>)

**Before taking MAVYRET, tell your healthcare provider about all of your medical conditions, including if you:**

- have ever had hepatitis B virus infection
- have liver problems other than hepatitis C virus infection.
- are pregnant or plan to become pregnant. It is not known if MAVYRET will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if MAVYRET passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you take MAVYRET.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

MAVYRET and other medicines may affect each other. This can cause you to have too much or not enough MAVYRET or other medicines in your body. This may affect the way MAVYRET or your other medicines work, or may cause side effects.

**Keep a list of your medicines to show your healthcare provider and pharmacist.**

- You can ask your healthcare provider or pharmacist for a list of medicines that interact with MAVYRET.
- **Do not start taking a new medicine without telling your healthcare provider.** Your healthcare provider can tell you if it is safe to take MAVYRET with other medicines.

**How should I take MAVYRET?**

- Take MAVYRET exactly as your healthcare provider tells you to take it. Do not change your dose unless your healthcare provider tells you to.
- Do not stop taking MAVYRET without first talking with your healthcare provider.
- Take 3 MAVYRET tablets at one time each day.
- Take MAVYRET with food.
- It is important that you do not miss or skip doses of MAVYRET during treatment.
- If you miss a dose of MAVYRET and it is:
  - Less than 18 hours from the time you usually take MAVYRET, take the missed dose with food as soon as possible. Then take your next dose at your usual time.
  - More than 18 hours from the time you usually take MAVYRET, do not take the missed dose. Take your next dose as usual with food.

If you take too much MAVYRET, call your healthcare provider or go to the nearest hospital emergency room right away.

**What are the possible side effects of MAVYRET?**

**MAVYRET can cause serious side effects, including:**

- **Hepatitis B virus reactivation.** See “What is the most important information I should know about MAVYRET?”

The most common side effects of MAVYRET include headache and tiredness.

These are not all the possible side effects of MAVYRET.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How should I store MAVYRET?**

- Store MAVYRET at or below 86°F (30°C).
- Keep MAVYRET in its original blister package until you are ready to take it.

**Keep MAVYRET and all medicines out of the reach of children.**

**General information about the safe and effective use of MAVYRET**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use MAVYRET for a condition for which it was not prescribed. Do not give MAVYRET to other people, even if they have the same symptoms that you have. It may harm them. You can ask your healthcare provider or pharmacist for information about MAVYRET that is written for health professionals.

**What are the ingredients in MAVYRET?**

**Active ingredients:** glecaprevir and pibrentasvir

**Inactive ingredients:** colloidal silicon dioxide, copovidone (type K 28), croscarmellose sodium, hypromellose 2910, iron oxide red, lactose monohydrate, polyethylene glycol 3350, propylene glycol monocaprylate (type II), sodium stearyl fumarate, titanium dioxide, and vitamin E (tocopherol) polyethylene glycol succinate. The tablets do not contain gluten.

Manufactured by AbbVie Inc., North Chicago, IL 60064.

MAVYRET is a trademark of AbbVie Inc. All other brands listed are trademarks of their respective owners and are not trademarks of AbbVie Inc. The makers of these brands are not affiliated with and do not endorse AbbVie Inc. or its products.

For more information go to [www.MAVYRET.com](http://www.MAVYRET.com) or call 1-800-633-9110.

This Patient Information has been approved by the U.S. Food and Drug Administration.  
03-B632

Issued: August 2017