

PATIENT INFORMATION
JUVMO™ (jooov'-moh)
(tavapadon)
tablets, for oral use

What is JUVMO?

JUVMO is a prescription medicine used for the treatment of Parkinson's disease (PD) in adults.

It is not known if JUVMO is safe and effective in children.

Before taking JUVMO, tell your healthcare provider about all your medical conditions, including if you:

- have low blood pressure (hypotension), or if your blood pressure decreases when standing (orthostatic hypotension).
- have or have had unusual urges such as problems with gambling, compulsive eating, compulsive shopping, and increased sex drive.
- have a mental health problem, such as psychosis.
- have seen or have been seeing things that are not there, hearing sounds, or feeling sensations that are not real (hallucinations).
- have trouble controlling your muscles (dyskinesia).
- are pregnant or plan to become pregnant. It is not known if JUVMO will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if JUVMO passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you take JUVMO.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicine, vitamins, and herbal supplements.

JUVMO and certain other medicines may affect each other and how they work.

Know the medicines you take. Keep a list of them and show your healthcare provider and pharmacist when you get a new medicine.

How should I take JUVMO?

- Take JUVMO exactly as your healthcare provider tells you to.
- Your healthcare provider may change your dose if needed.
- Take JUVMO by mouth with or without food.
- If you miss a dose of JUVMO, take it as soon as you remember on the same day, but do not take 2 doses at one time. If you do not remember until the following day, skip the missed dose and take JUVMO at your regular time.
- If you take too much JUVMO, call your healthcare provider or go to the nearest hospital emergency room right away.

What should I avoid while taking JUVMO?

- Do not change your body position too fast. Get up slowly from sitting or lying down. JUVMO can lower your blood pressure, which can cause nausea, sweating, dizziness, falling, or fainting when you stand up too quickly.

What are the possible side effects of JUVMO?

JUVMO may cause serious side effects, including:

- **low blood pressure. This is a common side effect and may be serious.** JUVMO can lower your blood pressure and make you feel dizzy, especially when you stand up (orthostatic hypotension). To help avoid this, do not get up

too fast from sitting or after lying down. You may be more likely to feel dizzy after starting JUVMO or when your dose is increased.

- **unusual urges.** Some people taking certain medicines to treat PD, including JUVMO, have had unusual urges. This may include gambling, compulsive eating, compulsive shopping, and increased sex drive. If you or your family members notice that you are having unusual urges or behaviors, talk to your healthcare provider.
- **seeing things that are not there, hearing sounds, or feeling sensations that are not real (hallucinations).** **This is a common side effect and may be serious.** Hallucinations can happen in people who use JUVMO. Tell your healthcare provider if you have hallucinations.
- **uncontrolled sudden movements (dyskinesia).** **This is a common side effect and may be serious.** If you have new dyskinesia, or your dyskinesia gets worse, tell your healthcare provider. This may be a sign that your dose of JUVMO or other medicines to control your PD may need to be adjusted.

The most common side effects of JUVMO include:

- nausea
- headache
- change in taste
- feeling tired (fatigue)
- vomiting
- dry mouth
- anxiety

Tell your healthcare provider if you have any side effect that bothers you or does not go away.

These are not all the possible side effects of JUVMO. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store JUVMO?

- Store JUVMO at room temperature between 68°F to 77°F (20°C to 25°C).

Keep JUVMO and all medicines out of the reach of children.

General information about the safe and effective use of JUVMO.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use JUVMO for a condition for which it was not prescribed. Do not give JUVMO to other people, even if they have the same symptoms that you have. It may harm them.

You can ask your pharmacist or healthcare provider for information about JUVMO that is written for health professionals.

What are the ingredients in JUVMO?

Active ingredient: tavapadon

Inactive ingredients:

0.25 mg and 1 mg tablets contain: lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, red iron oxide, sodium starch glycolate, talc, titanium dioxide, and yellow iron oxide.

5 mg, 10 mg, and 15 mg tablets contain: carmine, FD&C Blue No. 2, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, sodium starch glycolate, talc and titanium dioxide.

Marketed by: AbbVie Inc., North Chicago, IL 60064 U.S.A.

©2026 AbbVie. All rights reserved.

JUVMO and its design are trademarks of Cerevel Therapeutics, LLC, an AbbVie company.

For more information, call 1-888-817-4847 or go to www.JUVMO.com.